

# Biomarker 11: *Stensioeina granulata levis* FDO

From Earthwise

[Jump to navigation](#) [Jump to search](#)

This biomarker occurs in the Lamplugh Formation (early Coniacian, *M. cortestudinarium* Zone). The species is extremely rare in Britain, but it has been recorded in the lower *M. cortestudinarium* Zone (Bailey et al., 1984), and in Germany its range is given as "Ober Turon" (Koch, 1977).

**Correlation:** This biomarker can be placed in the BGS foraminiferal Biozone 13 (Wilkinson, 2000).

Retrieved from

'[http://earthwise.bgs.ac.uk/index.php?title=Biomarker\\_11:\\_Stensioeina\\_granulata\\_levis\\_FDO&oldid=1808](http://earthwise.bgs.ac.uk/index.php?title=Biomarker_11:_Stensioeina_granulata_levis_FDO&oldid=1808)'

[Category:](#)

- [Offshore foraminiferal biomarkers in the Chalk Group](#)

## Navigation menu

### Personal tools

- Not logged in
- [Talk](#)
- [Contributions](#)
- [Log in](#)
- [Request account](#)

### Namespaces

- [Page](#)
- [Discussion](#)

### Variants

### Views

- [Read](#)
- [Edit](#)
- [View history](#)
- [PDF Export](#)

## More

## Search

## Navigation

- [Main page](#)
- [Recent changes](#)
- [Random page](#)
- [Help about MediaWiki](#)

## Tools

- [What links here](#)
- [Related changes](#)
- [Special pages](#)
- [Permanent link](#)
- [Page information](#)
- [Cite this page](#)
- [Browse properties](#)

- This page was last modified on 1 October 2013, at 10:56.

- [Privacy policy](#)
- [About Earthwise](#)
- [Disclaimers](#)

