

# Chalk Biozone 12 - Callaiosphaeridium asymmetricum

From Earthwise

[Jump to navigation](#) [Jump to search](#)

**Age:** Early Campanian.

**Macrofaunal correlation:** The lowermost *O. pilula* Zone to the middle *G. quadrata* Zone.

**Definition:** The interval between the LAD of *Cassiculosphaeridia reticulata* to the LAD of *Callaiosphaeridium asymmetricum*.

**Comments:** The index for the top of this biozone, *Callaiosphaeridium asymmetricum*, has a well established LAD within the *Q. quadrata* Zone (Haq *et al.*, 1987; Costa & Davey, 1992).

**Lithostratigraphical correlation:** The Newhaven Chalk and Culver Chalk formations (Southern Province), the Flamborough Chalk Formation (Northern Province) and the Mackerel Formation (North Sea).

Retrieved from

'[http://earthwise.bgs.ac.uk/index.php?title=Chalk\\_Biozone\\_12\\_-\\_Callaiosphaeridium\\_asymmetricum&oldid=1565](http://earthwise.bgs.ac.uk/index.php?title=Chalk_Biozone_12_-_Callaiosphaeridium_asymmetricum&oldid=1565)'

[Category:](#)

- [Palynological zonation](#)

## Navigation menu

### Personal tools

- Not logged in
- [Talk](#)
- [Contributions](#)
- [Log in](#)
- [Request account](#)

### Namespaces

- [Page](#)
- [Discussion](#)

### Variants

## Views

- [Read](#)
- [Edit](#)
- [View history](#)
- [PDF Export](#)

## More

## Search

## Navigation

- [Main page](#)
- [Recent changes](#)
- [Random page](#)
- [Help about MediaWiki](#)

## Tools

- [What links here](#)
- [Related changes](#)
- [Special pages](#)
- [Permanent link](#)
- [Page information](#)
- [Cite this page](#)
- [Browse properties](#)

• This page was last modified on 30 September 2013, at 10:13.

- [Privacy policy](#)
- [About Earthwise](#)
- [Disclaimers](#)

