

Chalk Biozone 9 - *Endoscrinium campanula*

From Earthwise

[Jump to navigation](#) [Jump to search](#)

Age: Mid Coniacian to Early Santonian.

Macrofaunal correlation: The *M. coranguinum* Zone.

Definition: The interval between the LAD of *Florentinia deanei* to the LAD of *Endoscrinium campanula*.

Comments: The distinctive cavate species *Endoscrinium campanula* has a well established range top at the top of the *M. coranguinum* Zone (Foucher, 1979; Costa & Davey, 1992). A possible alternative index for the top of this biozone is the FAD of *Senoniasphaera protrusa* (see Foucher, 1979).

Lithostratigraphical correlation: The Seaford Chalk Formation (Southern Province), the Burnham Chalk and the lower Flamborough Chalk formations (Northern Province) and the Mackerel Formation (North Sea).

Retrieved from

'http://earthwise.bgs.ac.uk/index.php?title=Chalk_Biozone_9_-_Endoscrinium_camarula&oldid=1558'

Category:

- [Palynological zonation](#)

Navigation menu

Personal tools

- Not logged in
- [Talk](#)
- [Contributions](#)
- [Log in](#)
- [Request account](#)

Namespaces

- [Page](#)
- [Discussion](#)



Variants

Views

- [Read](#)
- [Edit](#)
- [View history](#)
- [PDF Export](#)

□

More

Search

Navigation

- [Main page](#)
- [Recent changes](#)
- [Random page](#)
- [Help about MediaWiki](#)

Tools

- [What links here](#)
- [Related changes](#)
- [Special pages](#)
- [Permanent link](#)
- [Page information](#)
- [Cite this page](#)
- [Browse properties](#)

- This page was last modified on 30 September 2013, at 10:09.

- [Privacy policy](#)
- [About Earthwise](#)
- [Disclaimers](#)



•



•



•