

Hydrogeology of Wales: Ordovician and Silurian aquifers - groundwater chemistry

From Earthwise

[Jump to navigation](#) [Jump to search](#)

This page is part of a category of pages that provides an updated review of the occurrence of groundwater throughout Wales.

Author(s): N S Robins and J Davies, British Geological Survey

Contributor(s): D A Jones, Natural Resources Wales and G Farr, British Geological Survey

The chemical composition of the groundwater in the Ordovician and Silurian strata is variable. This reflects the range of residence times as well as pH and redox. The rocks are dominated by poorly reactive minerals such as illite, chlorite and quartz, and the groundwaters tend to be weakly mineralised ([Shand et al., 2005](#)). The presence, locally, of sulphide and carbonate minerals form an important control on the evolution of the water chemistry. However, the largest variations are likely to occur with depth and boreholes that abstract waters from a variety of fractures at different depths produce waters of contrasting chemistry. The groundwaters are young, varying in age from weeks/months to a few decades.

The chemistry of the groundwater varies from very dilute waters dominated by Na, Cl and SO₄ (reflecting an important atmospheric input) to Ca-HCO₃ and mixed types. They vary from acidic to alkaline (pH 4.9 to 8.8) and from oxidising to reducing (Eh-79 to 514 mV). The pH is largely controlled by the degree of water-rock interaction, sulphide oxidation (producing acidity) and carbonate dissolution (producing alkalinity). The dominant process in most groundwaters is silicate dissolution, due to the lack of carbonate minerals present in the bedrock. In the western part of the Teifi Valley, the occurrence of carbonate shelly debris and calcite in the Irish Sea-derived drift has had a marked influence on the concentrations of Ca, Mg, HCO₃ and Sr.

Hydrogeology of Wales - contents

[Summary](#)

[Acknowledgements](#)

[Introduction](#)

[Geology and Groundwater](#)

[Topography, climate, land use and natural resources](#)

[Groundwater regulation](#)

[Issues](#)

[Precambrian and Cambrian](#)

[Groundwater occurrence in the Precambrian and Monian Supergroup](#)

[Groundwater occurrence in the Cambrian](#)

[Ordovician and Silurian](#)

[Groundwater occurrences](#)

[Groundwater studies](#)

Groundwater chemistry

[The Old Red Sandstone](#)

[Groundwater occurrences](#)

[Groundwater chemistry](#)

[Carboniferous](#)

[Carboniferous Limestone](#)

[Marros Group](#)

[Modelling the South Wales Coalfield](#)

[Coal Measures facies](#)

[Groundwater quality in the South Wales Coalfield](#)

[Permo-Triassic and Jurassic](#)

[Vale of Clwyd](#)

[Cheshire Basin, Dee catchment](#)

[South Wales](#)

[Quaternary aquifers](#)

[Groundwater occurrences](#)

[Afon Teifi](#)

[Upper Lugg catchment](#)

[Afon Cynffig coastal plain](#)

[Whiteford Sands](#)

[Newborough Warren](#)

[Management and regulation of groundwater](#)

[Groundwater abstraction](#)

[Need for management](#)

[Groundwater pollution](#)

[Management tools and future issues](#)

[References](#)

Retrieved from

'http://earthwise.bgs.ac.uk/index.php?title=Hydrogeology_of_Wales:_Ordovician_and_Silurian_aquifers_-_groundwater_chemistry&oldid=25641'

[Category](#):

- [Hydrogeology of Wales](#)

Navigation menu

Personal tools

- Not logged in
- [Talk](#)
- [Contributions](#)
- [Log in](#)
- [Request account](#)

Namespaces

- [Page](#)
- [Discussion](#)

Variants

Views

- [Read](#)
- [View source](#)
- [View history](#)
- [PDF Export](#)

□

More

Search

Navigation

- [Main page](#)
- [Recent changes](#)
- [Random page](#)
- [Help about MediaWiki](#)

Tools

- [What links here](#)
- [Related changes](#)
- [Special pages](#)
- [Permanent link](#)
- [Page information](#)
- [Cite this page](#)
- [Browse properties](#)

• This page was last modified on 18 February 2016, at 09:31.

- [Privacy policy](#)
- [About Earthwise](#)
- [Disclaimers](#)

