

Subprionocyclus normalis Subzone

From Earthwise

[Jump to navigation](#) [Jump to search](#)

The subzone is characterised by *Subprionocyclus normalis* and *Baculites undulatus*, and has been proved in Hertfordshire, Bedfordshire and Surrey (Gale, 1996). The subzone appears to equate with the upper *S. plana* Zone of the standard macrofossil zonal scheme (Gale, 1996).

Retrieved from

'http://earthwise.bgs.ac.uk/index.php?title=Subprionocyclus_normalis_Subzone&oldid=1607'

Category:

- [International ammonite zonation for the Turonian & Early Coniacian](#)

Navigation menu

Personal tools

- Not logged in
- [Talk](#)
- [Contributions](#)
- [Log in](#)
- [Request account](#)

Namespaces

- [Page](#)
- [Discussion](#)

Variants

Views

- [Read](#)
- [Edit](#)
- [View history](#)
- [PDF Export](#)

More

Search

Navigation

- [Main page](#)
- [Recent changes](#)
- [Random page](#)
- [Help about MediaWiki](#)

Tools

- [What links here](#)
- [Related changes](#)
- [Special pages](#)
- [Permanent link](#)
- [Page information](#)
- [Cite this page](#)
- [Browse properties](#)

• This page was last modified on 30 September 2013, at 11:54.

- [Privacy policy](#)
- [About Earthwise](#)
- [Disclaimers](#)

